

We need a new GSSP for the base of the Jurassic System

Spencer G. LUCAS¹, Karl KRAINER², Lawrence H. TANNER³, David G. TAYLOR⁴

Since 1961, the IUGS International Commission on Stratigraphy (ICS) has been developing and standardizing the chronostratigraphic scale by defining chronostratigraphic boundaries by identifying Global Stratotype Sections and Points (GSSPs). To achieve this, the chronostratigraphic divisions of each geological system are being developed by a Subcommittee, and each subcommittee creates working (or task) groups to identify candidate sections and criteria for the selection of GSSPs defining the bases of relevant stage units.

In 1984, the Jurassic Subcommittee created a working group to find a GSSP for the base of the Hettangian Stage, which is the lowest Stage of the Jurassic, though efforts to define a Jurassic base had begun in the 1960s (Maubeuge, 1964). After about 30 years of deliberation, the working group had identified four GSSP candidates: (1) St. Audries Bay, Somerset, UK; (2) Kennecott Point, Kunga Island, British Columbia, Canada; (3) New York Canyon area, Nevada, USA; and (4) the Utcabamba Valley, Peru (Fig. 1). It had long been known that a major turnover of the Ammonoidea characterizes the Triassic–Jurassic transition, so many workers favored an ammonoid signal to define the Jurassic base, though other signals, including radiolarian turnover and a carbon isotope excursion had their advocates (Fig. 2) (for a review, see Warrington *et al.*, 1994; Lucas *et al.*, 2007; Hillebrandt *et al.*, 2013).

In 2007, two new sections were added to the roster of potential GSSP candidates: Waterloo Bay, Northern Ireland, and Kuhjoch, Austria (Fig. 1). Furthermore, the group to vote on a GSSP was expanded to an unprecedented 75 members, 47 of them from Europe. The section at Waterloo Bay had been studied and published on since the 1800s, but nothing had been published on the newly discovered Kuhjoch section prior to its proposal by Hillebrandt *et al.* (2007) as a GSSP candidate (published in the ISJS Newsletter).

Despite this, the vote of the working group took place in February–April 2008, and the Kuhjoch section was chosen as the GSSP for the base of the Hettangian, and the Nevada section as an Auxiliary Stratotype Section and Point. The Jurassic Subcommittee approved the decision in June 2008, the International Committee on Stratigraphy did so in May 2009 and, in April 2010, it was ratified by the Executive Committee of the IUGS. Hillebrandt *et al.* (2013) presented a detailed description of the Kuhjoch GSSP. Since its initial proposal, only one article (Hillebrandt, Krystyn, 2009) was published on the Kuhjoch section before it was ratified as the GSSP. Thus, the selection of Kuhjoch as the GSSP for the base of the Hettangian was added at the eleventh hour to a list of long studied and well understood candidates. Since nothing had been published on the Kuhjoch section prior to its proposal, its evaluation and discussion from the expanded group of voting members prior to its ratification was minimal.

Although the Kuhjoch section appears to contain a satisfactory paleontological and geochemical record, there are other aspects of the section that should have disqualified it from consideration as a GSSP. Palotai *et al.* (2017) recently restudied the Kuhjoch section, demonstrating that it is extensively disturbed tectonically – all the incompetent beds are foliated, there are tight to isoclinal folds in the strata and a reverse fault cuts through the GSSP section, so that part of the section is

¹ New Mexico Museum of Natural History, 1801 Mountain Road NW, Albuquerque, New Mexico 87104 USA; spencer.lucas@state.nm.us.

² Institute of Geology, University of Innsbruck, Innrain 52, A-6020 Innsbruck, Austria; karl.krainer@uibk.ac.at.

³ Department of Biology, LeMoyne College, Syracuse, New York 13214 USA; tannerlh@lemoyne.ed.

⁴ Department of Geology, Portland State University, 1721 SW Broadway Ave., Portland OR 97201 and Department of Earth Sciences, 1272 University of Oregon, Eugene OR 97403 USA; Blitz124@comcast.net.





Fig. 1. GSSP candidates for the base Jurassic (Hettangian) GSSP voted on in 2008 (modified from Lucas *et al.*, 2007)

tectonically omitted. Hillebrandt *et al.* (2013, table 1) stated that at Kuhjoch there was “an absence of synsedimentary and tectonic disturbance near [the] boundary level”. However, Palotai *et al.* (2017: 2475) concluded, “the Kuhjoch sections do not fulfill the specific requirement for a GSSP regarding the absence of tectonic disturbances near boundary level”. Indeed, such tectonic disturbance of the section raises serious questions about the stratigraphic position and relationships of the biostratigraphic and chemostratigraphic events in the Kuhjoch section depicted by Hillebrandt *et al.* (2007, 2013). Additionally, the strata of the Kuhjoch sections are heavily weathered; sampling for paleontological or geochemical analysis requires excavation with heavy equipment to expose fresh beds. Hence, each successive study examines slightly different rocks than previous studies (*e.g.*, Tanner *et al.*, 2016).

The choice of the Kuhjoch GSSP is a cautionary tale for those seeking GSSP-based chronostratigraphic definitions (also see Lucas, 2018). Typically, the process of choosing a GSSP takes at least a decade, as the relevant working group carefully evaluates and chooses a section that is very well understood and meets as many of the ICS criteria for a GSSP as possible. Furthermore, candidate sections are almost always sections that have been long known, long studied and have adequate published documentation. Kuhjoch was not such a section, and though it fell from the sky in 2007, it was also not a *deus ex machina*.

Clearly, a new GSSP for the base of the Jurassic is needed. The Jurassic Subcommittee should create a new working group for that purpose. A possible solution could be to simply designate the auxiliary GSSP section in the New York Canyon area, Nevada, as the GSSP. That section continues to be the focus of much research (*e.g.*, Ritterbush *et al.*, 2014; Hodges, Stanley, 2015; Thibodeau *et al.*, 2016) and will serve as a GSSP without the tectonic complications that plague the current base-Hettangian GSSP at Kuhjoch, Austria.

Fig. 2. Succession of potential primary signals for the base-Jurassic GSSP discussed and voted on in 2008 (modified from Lucas *et al.*, 2007). The green interval encompasses the range of most of the possible GSSP levels for the base of the Hettangian



REFERENCES

- HILLEBRANDT A. von., KRYSSTYN L., 2009 – On the oldest Jurassic ammonites of Europe (Northern Calcareous Alps) and their global significance. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, **253**: 163–195.
- HILLEBRANDT A. von., KÜRSCHNER W.M., KRYSSTYN L., 2007 – A candidate GSSP for the base of the Jurassic in the Northern Calcareous Alps (Kuhjoch section, Karwendel Mountains, Tyrol, Austria). *International Subcommission for Jurassic Stratigraphy Newsletter*, **34**: 2–20.
- HILLEBRANDT A. von., KRYSSTYN L., KÜRSCHNER W.M., BONIS N.R., RUHL M., RICHOZ S., SCHOBEN M.A.N., ULRICH M., BOWN P.R., KMENT K., MCROBERTS C.A., SIMMS M., TOMÁŠOVÝCH A., 2013 – The global stratotype sections and point (GSSP) for the base of the Jurassic System at Kuhjoch (Karwendel Mountains, Northern Calcareous Alps, Tyrol, Austria). *Episodes*, **36**: 162–198.
- HODGES M.S., STANLEY JR, G.D., 2015 – North American coral recovery after the end-Triassic mass extinction, New York Canyon, Nevada, USA. *GSA Today*, **25**: 4–9.
- LUCAS S.G., 2018 – The GSSP method of chronostratigraphy: A critical review. *Frontiers in Earth Science*, **6**, 191: 1–18.
- LUCAS S.G., TAYLOR D.G., GUEX J., TANNER L.H., KRAINER K., 2007 – The proposed global stratotype section and point for the base of the Jurassic System in the New York Canyon area, Nevada, USA. *New Mexico Museum of Natural History and Science Bulletin*, **40**: 139–168.
- MAUBEUGE P.-L., 1964 – Résolutions du Colloque. In: Colloque du Jurassique à Luxembourg 1962 (Ed. P.-L. Mauberge). *Ministère des Arts et de Sciences, Publications de l'Institut Grand-Ducal, Section des Sciences Naturelles, Physiques et Mathématiques*: 77–80.
- PALOTAI M., PÁLFY J., SASVÁRI Á., 2017 – Structural complexity at and around the Triassic–Jurassic GSSP at Kuhjoch, Northern Calcareous Alps, Austria. *International Journal of Earth Science*, **106**: 2475–2487.
- RITTERBUSH K.A., BOTTJER D.J., CORSETTI F.A., ROSAS S., 2014 – New evidence on the role of siliceous sponges in ecology and sedimentary facies development in eastern Panthalassa following the Triassic–Jurassic mass extinction. *Palaios*, **29**: 652–668.
- TANNER L.H., KYTE F.T., RICHOZ S., KRYSSTYN L., 2016 – Distribution of iridium and associated geochemistry across the Triassic–Jurassic boundary in sections at Kuhjoch and Kendlbach, Northern Calcareous Alps, Austria. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **449**: 13–26.
- THIBODEAU A.M., RITTERBUSH K., YAGER J.A., WEST A.J., IBARRA Y., BOTTJER D.J., BERELSON W.M., BERGQUIST B.A., CORSETTI F.A., 2016 – Mercury anomalies and the timing of biotic recovery following the end-Triassic mass extinction. *Nature Communications*, **7**: 11147.
- WARRINGTON G., COPE J.C.W., IVIMEY-COOK H.C., 1994 – St Audrie's Bay, Somerset, England: a candidate Global Stratotype Section and Point for the base of the Jurassic System. *Geological Magazine*, **131**: 191–200.

